

\*ProBios tube - Equine formula (enteritis issues-diarrhea or ABX admin

**GIVE ANY TIME YOU GIVE ABX)**

\*Liquid Imodium – (for diarrhea-HELPS STOP 1-2 CC'S depending on weight)

\*GasX strips or simethicone drops (gas issues with enteritis BLOAT 1-2 DROPPERS FULL)

Ivermectin or Ivomec –injectable type 1% .5cc for 10# rabbit administer topically)

\*Tums-fruit flavored (calcium boost for kindling does-helps with labor(3-4-crush and give with water via syringe or eye dropper if needed; GIVE 1-2 EVERY DAY FOR 3-4 DAYS AFTER KINDLING)

\*Injectable Penicillin **never straight penicillin**: I use PenG w/Procaine (mastitis) – for dosing see [www.medirabbit.com](http://www.medirabbit.com)

\*Triple antibiotic ointment, no pain meds in it (general wounds)

Papaya Enzyme – Molting – helps avoid fur block

\*Vetricyn – for all wounds, eye issues, everything topical

\*Terramycin eye ointment or Veterycin spray (nestbox eye)

Anti-fungal cream – for fungal infection (usually hocks)

\*Disposable Scalpels-- rounded and pointed blades (lancing abscesses)

\*Vet wrap, sterile gauze pads, telfa pads or band-aides (FOR YOU!),

\*Syringes- feeding type, tuberculin (1cc) and larger, 6-12 CC sizes

\*Hypodermic needles..18g to draw with, 22g to inject with

\*Blue kote, Bag Balm (for wounds, ie. sore hocks, etc.)

\*Spray Iodine, different dilutions (general cleaning, wound care)

\*Vinegar and rubbing alcohol (general cleaning); Bleach & Ammonia

\*Hay – when off feed or diarrhea; or molting

\*Oats – when off feed

Flour or corn starch – to stop bleeding when clipping nails (oops)

\*Scissors (we have bandage scissors, and small, straight scissors)

\*Tweezers- sharp and blunt

KEVLAR GLOVES-SAVE YOURSELF

Eye Droppers – for administering liquids or meds

\*Scale (charting weight); \*Calculator (for calculating dosages for ABX)

\*Dosage charts (for your commonly used meds)

Shannon Neal Pankow - Carla's changes

ABX= antibiotics